

DIAGRAMMATIC THEORY OF THE ANDERSON IMPURITY MODEL WITH FINITE COULOMB INTERACTION

K. Haule^{a,b}, S. Kirchner^b, H. Kroha^b, P. Wölfle^b

^a *Institute Josef Stefan, Department of Theoretical Physics (F1)*

SI-1000 Ljubljana, Slovenia

^b *Institut für Theorie der Kondensierten Materie, Universität Karlsruhe*

D-76128 Karlsruhe, Germany

Abstract We developed a self-consistent conserving pseudo particle approximation for the Anderson impurity model with finite Coulomb interaction, derivable from a Luttinger Ward functional. It contains an infinite series of skeleton diagrams built out of fully renormalized Green's functions. The choice of diagrams is motivated by the Schrieffer Wolff transformation which shows that singly and doubly occupied states should appear in all bare diagrams symmetrically. Our numerical results for T_K are in excellent agreement with the exact values known from the Bethe ansatz solution. The low energy physics of non-Fermi liquid Anderson impurity systems is correctly described while the present approximation fails to describe Fermi liquid systems, since some important coherent spin flip and charge transfer processes are not yet included. It is believed that CTMA (Conserving T-matrix approximation) diagrams will recover also Fermi liquid behavior for Anderson models with finite Coulomb interaction as they do for infinite Coulomb interaction.

1. INTRODUCTION

Anderson impurity models have been of considerable interest recently for studying the transport in mesoscopic structures [1] as well as for describing strong correlations on the lattice in the limit of large dimensions [2]. In both cases the doubly occupied impurity site, which is characterized by an additional energy cost U , is essential for the correct physical description.

In terms of pseudofermion (f_σ) and slave boson operators (a, b) the Anderson model is defined by the Hamiltonian

$$H = \sum_{\vec{k}, \sigma} \epsilon_{\vec{k}} c_{\vec{k}\sigma}^\dagger c_{\vec{k}\sigma} + \epsilon_d \left(\sum_{\sigma} f_{\sigma}^\dagger f_{\sigma} + 2a^\dagger a \right) +$$

$$+ \sum_{\vec{k}, \sigma} V_k (c_{\vec{k}\sigma}^{\dagger} b^{\dagger} f_{\sigma} + \sigma c_{\vec{k}\sigma}^{\dagger} f_{-\sigma}^{\dagger} a + h.c.) + U a^{\dagger} a;$$

where ϵ_d is the energy of the local d level, U is the Coulomb interaction and $V_{\vec{k}}$ are the hybridization matrix elements. A physical electron in the local level is created with the operator $d_{\sigma} = b^{\dagger} f_{\sigma} + \sigma f_{-\sigma}^{\dagger} a$, where b represents the empty level and a the doubly occupied one (in the following we will call a and b the heavy and the light boson, respectively). For the representation to be exact, the local operator constraint $Q \equiv \sum_{\sigma} f_{\sigma}^{\dagger} f_{\sigma} + b^{\dagger} b + a^{\dagger} a = 1$ must be fulfilled at all times. This is guaranteed by a suitable projection technique [4].

2. METHOD

It is well known that the Non-Crossing Approximation (NCA) gives the correct Kondo temperature T_K in the limit of large Coulomb repulsion U while it fails to recover the correct Kondo temperature for finite U . The reason may be seen from the Schrieffer-Wolff transformation [3] where the effective coupling constant has two equally important terms for $U \simeq -2\epsilon_d > 0$

$$J = -\frac{V^2}{\epsilon_d} + \frac{V^2}{\epsilon_d + U} \quad (1.1)$$

where just one of them survives in the large U limit. In Eq. (1.1) the first (second) term is generated by exchange of a bare light (heavy) boson. This implies that the light (b) and heavy (a) bosons should appear symmetrically in all bare diagrams. In order to generate the corresponding diagrams of the Luttinger-Ward functional one may start with the NCA diagram for the $U \rightarrow \infty$ case (first diagram in Fig. 1a)) and replace the light boson line by a heavy boson, exchanging the conduction electron lines appropriately. The result is the second diagram. This is not sufficient, however, because the diagrams should be symmetric in a, b on the level of bare diagrams. One may show that this exchange generates an infinite series of diagrams, each containing a circle of pseudoparticle lines with one light boson and n heavy bosons (or vice versa) dressed by conduction electron lines spanning at most three pseudoparticle lines (see Fig. 1a)). The corresponding self-energies are obtained by functional differentiation and are shown in Fig. 1b) in terms of vertex functions (the dash- and cross-shaded half-circle areas).

The Green's function of the physical d electrons can be calculated from the exact relation that can be derived with the help of the equation of motion

$$G_{\vec{k}\vec{k}'}(\omega) = \delta_{\vec{k}\vec{k}'} G_{\vec{k}\vec{k}}^0(\omega) + \sum_{\vec{k}\vec{k}'} G_{\vec{k}\vec{k}'}^0(\omega) V_{\vec{k}} G_d(\omega) V_{\vec{k}'}^* G_{\vec{k}'\vec{k}}^0(\omega). \quad (1.2)$$




Figure 1 a) The Luttinger Ward functional, b) the Bethe-Salpeter equations for the T-matrix vertices (dark and light shaded areas), c) the Contributions to the irreducible self energies for pseudo-particles as well as for conduction electrons are shown.

The local conduction electron Green's function can be expressed in terms of the local conduction electron self energy Σ_c (see Fig. 1c)),

$$G_d(\omega) = \frac{1}{V^2} \Sigma_c(\omega). \quad (1.3)$$

3. NUMERICAL RESULTS

The Kondo temperature was extracted from the width of the Abrikosov-Suhl peak in the local electron Green's function. The values we get from the present approximation are in excellent agreement with the exact values known from the Bethe ansatz solution, as can be seen in Fig. 2. The simple Non-Crossing Approximation (the first two terms in Φ in Fig. 1a) gives Kondo temperatures several orders of magnitude too low. Including one additional diagram does not increase T_K sufficiently (*UNCA*) (here we disagree with the corresponding claim in Ref. [5]). An *infinite* series of *skeleton* ladder diagrams is needed to get the correct scaling of T_K with the Coulomb energy U and the local electron energy ϵ_d .




Figure 2 Left: Kondo temperature for various parameters (E_d and U). Solid lines are values of T_K from the Bethe-ansatz solutions. Right: Local electron spectral function calculated with NCA, UNCA and the ladder approximation. The width of the peak is proportional to the Kondo temperature.

4. SUMMARY

In summary, we have shown that an infinite number of skeleton diagrams is necessary to recover the correct dynamic energy scale (T_K) in the Anderson impurity model with finite Coulomb interaction energy. We found a systematic way to obtain those diagrams and have shown that the approximation is derivable from Luttinger-Ward functional and therefore conserving. Numerical results show excellent agreement with the exact values of T_K .

This work was funded by the ESF Program on "Fermi-liquid instabilities in correlated metals" (FERLIN), and by SFB 195 der Deutschen Forschungsgemeinschaft.

References

- [1] For an overview see *Mesoscopic Electron Transport* L. L. Sohn, L. P. Kowenhoven, G. Schön, NATO ASI Series E, Vol. 345 (Kluwer, 1997).
- [2] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996) .
- [3] J. R. Schrieffer in P. A. Wolff, Phys. Rev. **149**, 491 (1966).
- [4] J. Kroha, P. Wölfle, and T. A. Costi, Phys. Rev. Lett. **79**, 261 (1997); J. Kroha and P. Wölfle, Acta Phys. Pol. B **29**, (12) 3781 (1998), cond-mat/9811074; J. Kroha and P. Wölfle, Advances in Solid State Phys. **39**, 271 (1999).
- [5] Pruschke et al. Z. Phys. B **74**, 439-449 (1989).